

CAS BIOFINDER DISCOVERY PLATFORM™

快速使用指南 (QUICK GUIDE)

大綱

- CAS BioFinder簡介與界面介紹
- 配體檢索與配體詳情信息（配體與細節概述）
- 骨架檢索與構效關係分析（支架與搜救分析）
- 蛋白/通路/靶點、疾病與生物標誌物（Proteins and Pathways;疾病與生物標記）
- 生物活性預測（Predictive Analytics）

科學資訊管理對新葯研發至關重要

低增長時代的特點對創新的要求極大提升，而在當今的創新環境下，數據管理任務艱巨


- 信息鴻溝會導致創新管線薄弱、研發效率低下，以及意外的知識產權風險
- 由於數據量和數據類型日益複雜，藥物發現流程中，整合和分析關鍵的研發數據資訊，對於迅速做出關鍵決策至關重要。
- 持續的數據管理有助於開發差異化的成果，支撐創新藥物的發現，確保藥物研發成功進入審批階段，並可避免後續智慧財產權問題。

来源：CAS Insights洞察报告 [Challenges and opportunities in data for drug discovery | CAS](#)

生命科學領域研究離不開重要的數據資訊基礎， 但為什麼卻難以加以利用？

缺乏關聯性

生命科學研究和數據分散在各個資料庫、
期刊和專利等眾多互不關聯的資源之中



未一致化

不同資源對相同資訊的描述各不相同，
因此很難對數據進行分析和比較



數據問題導致延緩創新進程，增加財務風險

遺漏見解



這些見解能夠為研究人員制定研發戰略提供指導

僅在美國，每年就有5,000到6,000起專利侵權和專利無效的訴訟，法院判定的賠償金額接近 50 億美元¹。

耗費時間



生命科學數據量每 14 年翻一番²

數據科學家將 50%-80% 的時間都花費在收集和整理用於分析模型的數據資訊的瑣碎的工作中³

1. Vuleta, B. (2023). 25 Patent Litigation Statistics - High-Profile Feuds about Intellectual Property. LegalJobs. <https://legaljobs.io/blog/patent-litigation-statistics/>

2. Bornmann, L. et al. (2021). Growth rates of modern science: a latent piecewise growth curve approach to model publication numbers from established and new literature databases. (Humanities and Social Sciences Communications), 8, 224. <https://doi.org/10.1057/s41599-021-00903-w>

3. Lohr, S. (2014.) For Big-Data Scientists, 'Janitor Work' Is Key Hurdle to Insights. The New York Times. August 17, 2014. <https://www.nytimes.com/2014/08/18/technology/for-big-data-scientists-hurdle-to-insights-is-janitor-work.html>

4. Genetic Engineering & Biotechnology News Top 25 Biotech Companies of 2019. 5. Pharm Exec's Top 50 Companies 2020. 6. Shanghai Ranking's Global Ranking of Academic Subjects 2020.

利用CAS BioFinder 加速早期藥物研發



CAS BioFinder 檢索介面



CAS Newton Ligands Scaffolds Proteins **Diseases**

Search by protein, ligand, disease or draw a structure.



Draw



CAS Newton

Chat with CAS Newton to answer your drug discovery questions.



Predictive Analytics

Upload and analyze ligand data sets using algorithms to find bioactivity data insights.



Search Sequences **BETA**

Query BLAST algorithms for nucleotide and protein based sequences.

結構繪製面板

Recent Search History

[View Search History](#)

7 May 2026

近期檢索歷史

CAS Newton

Research

10:14 AM

雙特異抗體中靶點是pd1-vegfr的藥物

View Results



Search Complete

Edit search
Delete search

大綱

- CAS BioFinder簡介與檢界面介紹（介紹與介面）
- 配體檢索與配體詳情信息（配體與細節概述）
- 骨架檢索與構效關係分析（支架與搜救分析）
- 蛋白/通路/靶點、疾病與生物標誌物（Proteins and Pathways; 疾病與生物標記）
- 生物活性預測（Predictive Analytics）

基於靶點、疾病、藥物名稱以及結構，檢索資訊

Ligands

Scaffolds

BTK

IBTK

Tyrosine-protein kinase BTK

Tyrosine-protein kinase BTK/M

BTKh

BTK Orange II

BTK Black 10BX

BTK Victoria Blue

BTK deficiency

自動提示敘詞表

CAS BioFinder

Ligands BTK

×

Q

Draw

Filters

Chemical Filters

Rule of 5 Filter Preset

This will automatically be applied.

pValue

Modality

Druglikeness

ADME

Commercial Availability

Approval Status

Approval Authority

Biological Filters

Target

Target Type

Disease

Ligands search for BTK

Ligands Scaffolds Pharmacology

29,502 Results


Sort: Relevance

Predictive Analytics

1

936563-96-1

Ibrutinib




Metabolites: 3

Proteins: 687

2

1420477-60-6

4-[8-Amino-3-[(2S)-1-(1-oxo-2-butyn-1-yl)-2-pyrrolidinyl]imidazo[1,5-a]pyrazin-1...




Metabolites: 0

Proteins: 203

3

1691249-45-2

(7S)-4,5,6,7-Tetrahydro-7-[1-(1-oxo-2-propen-1-yl)-4-piperidinyl]-2-(4-phenoxyph...



Metabolites: 0

Proteins: 36

4

Relevance

CAS RN: Ascending

CAS RN: Descending

Number of Targets: Ascending

Number of Targets: Descending

Number of Metabolites: Ascending

Number of Metabolites: Descending

Suppliers

检索结果排序

5

2095280-64-9

N-[3-[1,6-Dihydro-1-methyl-5-[[4-(4-

Metabolites: 0

Proteins: 215

7

1803358-13-5

N-[2-Methyl-5-[2-oxo-9-(1H-pyrazol-1-

Metabolites: 0

Proteins: 215

10

© 2025 American Chemical Society. All rights reserved.

CAS
A division of the
American Chemical Society

篩選具有類似先導化合物特性的小分子配體

一鍵應用Rule of 5

Chemical Filters

☒ Rule of 5 Filter Preset
This will automatically be applied.

ADME

Partition Coefficient (logP)
-1.17 to 7.18
-1.17 to 5.00

Solubility (logS)
-8.94 to 0.58
-8.94 to 0.58

Blood-Brain Barrier Permeability (logBB)
-1.94 to 0.99
-1.94 to 0.99

Permeability Glycoprotein (Pgp)
0 to 1
0 to 1

Polar Surface Area (PSA) (Å²)
20 to 208
20 to 140

Plasma Protein Binding (PPB)
16 to 100
16 to 100

Modality

Druglikeness

Hydrogen Bond Acceptors
2 to 13
2 to 10

Hydrogen Bond Donors
0 to 7
0 to 5

Freely Rotatable Bonds
0 to 13
0 to 10

Molecular Weight (g/mol)
260 to 587
260 to 500

Molar Refractivity (m³/mol)
58 to 152
58 to 130

Number of Atoms
11 to 36
20 to 36

[Reset Filter](#) [Apply](#)

Ligands search for BTK

Ligands Scaffolds Pharmacology

Applied Filters (8) Hydrogen Bond Acceptors: 0 to 10 X Hydrogen Bond Donors: 0 to 5 X Freely Rotatable Bonds: 0 to 10 X Molecular Weight (g/mol): 0 to 500 X Molar Refractivity (m³/mol): 40 to 130 X Partition Coefficient (logP): up to 5 X Polar Surface Area (PSA) (Å²): 0 to 140 X

9,273 Results

1 [936563-96-1](#)
ibrutinib

Metabolites: 3 | Proteins: 687

2 [1202757-89-8](#)
CC 292

Metabolites: 7 | Proteins: 215

3 [1803358-13-5](#)
N-[2-Methyl-5-[2-oxo-9-(1H-pyrazol-1-yl)benzo[h]-1,6-naphthyridin-1(2H)-yl]phenyl]...

Metabolites: 7 | Proteins: 16

5 [1803358-31-7](#)
N-[2-Methyl-5-[2-oxo-9-(1H-pyrazol-1-yl)benzo[h]-1,6-naphthyridin-1(2H)-yl]phenyl]...

Metabolites: 8 | Proteins: 15

6 [1202755-92-7](#)
N-[3-[[5-Methyl-2-(phenylamino)-4-pyrimidinyl]amino]phenyl]-2-propenamide

Metabolites: 12 | Proteins: 12

7 [302962-49-8](#)
Dasatinib

Metabolites: 4 | Proteins: 733

9 [1415823-49-2](#)
1-[4-[[[6-Amino-5-(4-phenoxyphenyl)-4-pyrimidinyl]amino]methyl]-...

Metabolites: 12 | Proteins: 12

10 [1202755-89-2](#)
(2E)-4-(Dimethylamino)-N-[3-[5-fluoro-4-[(3-methylphenyl)amino]-...

Metabolites: 12 | Proteins: 12

11 [1202757-97-8](#)
1-[[[3R]-3-[[5-Fluoro-2-[(3-methoxyphenyl)amino]-4-pyrimidinyl]oxy]-...

Metabolites: 4 | Proteins: 733

- 氢键受体 < 10
- 氢键供体 < 5
- 自由旋转键 < 10
- 分子量 < 500
- 摩尔折射率：
40-130 m³/mol
- 原子数: 20-70
- logP < 5
- 极性表面积：
0-140 Å²

配體檢索結果的縱覽與聚焦

Disease

Enter a Disease... Search

- ☐ 0 Selected
- ☐ Rheumatoid arthritis
- ☐ Cancer
- ☐ Autoimmune disease
- ☐ Asthma
- ☐ Multiple sclerosis
- ☐ Myasthenia gravis
- ☐ Inflammatory bowel disease
- ☐ Crohn's disease
- ☐ Psoriasis
- ☐ Systemic lupus erythematosus
- ☐ Sjogren's syndrome
- ☐ Allergic disease
- ☐ Lupus erythematosus
- ☐ Graves' disease
- ☐ Osteoarthritis
- ☐ Leukemia
- ☐ Addison's disease
- ☐ Allergic rhinitis
- ☐ Lymphoma
- ☐ Type 1 diabetes mellitus
- ☐ B-cell lymphoma
- ☐ Multiple myeloma
- ☐ Chronic lymphocytic leukemia
- ☐ Scleroderma
- ☐ Diabetes mellitus

← Prev Next →

Source Filters

- Document Type
- Publication Year
- Language

Ligands search for BTK

Ligands Scaffolds Pharmacology

Applied Filters (1) Target: Tyrosine-protein kinase BTK X Clear all filters

29,502 Results

Filters

Chemical Filters

Rule of 5 Filter Preset Off This will automatically be applied.

pValue

Modality

- ☐ Small Molecule
- ☐ Protein/Peptide
- ☐ Other
- ☐ Coordination Compound
- ☐ Antibody
- ☐ Polymer
- ☐ Antibody Drug Conjugate

ADME

Commercial Availability

Approval Status

Approval Authority

Biological Filters

Target 靶点

Target Type 疾病

Disease

- ☐ Rheumatoid arthritis
- ☐ Cancer
- ☐ Autoimmune disease
- ☐ Asthma
- ☐ Multiple sclerosis

Organism

Parameter

Function

Experiment Type

- ☐ in vitro
- ☐ in vivo

View All Apply

有機分子功能

實驗類型

Approval Status

- ☐ Other
- ☐ Investigational
- ☐ US Approved
- ☐ Possibly Marketed Outside US
- ☐ Designated
- ☐ US Previously Marketed
- ☐ US Approved OTC

View Fewer Apply

Results

ID	Name	Metabolites	Proteins
936563-96-1	Ibrutinib	3	687
2095280-64-9	N-[3-{[1,6-Dihydro-1-methyl-5-[[4-(4-morpholinylcarbonyl)-3-[(1-oxo-2-propen-1-yl)...	15	13
2095280-65-0	N-[3-{[1,6-Dihydro-1-methyl-5-[[4-(4-morpholinylcarbonyl)-3-[(1-oxo-2-propen-1-yl)...	16	13
2101700-15-4	5-Amino-3-[4-[[[5-fluoro-2-methoxybenzoyl]amino]methyl]phenyl]-1-[(1S)-2,2-tri...	15	13
1803358-31-7	N-[2-Methyl-5-[2-oxo-9-(1H-pyrazol-1-yl)benzo[h]-1,6-naphthyridin-1(2H)-yl]phenyl...	16	13
1202755-92-7	N-[3-[[5-Methyl-2-(phenylamino)-4-pyrimidinyl]amino]phenyl]-2-propanamide	7	16

縱覽配體檢索結果集相關骨架結構與藥理學數據

查看配體檢索結果
關聯的骨架結構

Ligands search for BTK

Ligands **Scaffolds** Pharmacology

Applied Filters (1) Target: Tyrosine-protein kinase BTK X Clear all filters

10,870 Results

1

Ligands: 1 | Proteins: 1

2

Ligands: 1 | Proteins: 1

5

Ligands: 1 | Proteins: 1

6

Ligands: 1 | Proteins: 1

7

Ligands: 1 | Proteins: 1

8

Ligands: 1 | Proteins: 2



點擊View，可查看原文中的Assay的細節

探索配體的詳細資訊

Doxorubicin
CAS Registry Number: 23214-92-8

112K1,642RetrosynthesisPredictive Analytics

View Associated Scaffold

View in CAS SciFinder

連結到CAS SciFinder
查看關聯資訊

SummaryPharmacologyADMEToxicologyProteins (313)Diseases (301)Related Immunotherapeutics (12)Chemical SpaceMetabolites (4)Similar Ligands (764)

配體相關信息標籤頁

藥物情報資訊

CAS DRUG INTELLIGENCE

Approved Drug

Doxorubicin is an antineoplastic in the anthracycline class. General properties of drugs in this class include: interaction with DNA in a variety of different ways including intercalation (squeezing between the base pairs), DNA strand breakage and inhibition with the enzyme topoisomerase II. Most of...

View More

STATUS

Approval Status: US Approved
Approval Authority: US FDA
Approval Year: 1974, 2024
Originator: Farmitalia

INDICATIONS

Ovarian cancer
Modality: Primary
Highest Phase: Approved
[Source](#)

View All

PRIMARY TARGET

DNA topoisomerase II
Function: Inhibitor
Measurement: IC50 2.67 μM
[Source](#)

View All

View Related Drugs

配體屬性資訊

H bond acceptors*H bond donors*Molecular weight (g/mol)Molar refractivity (m²/mol)Number of atomsFreely rotatable bonds*logP*logSlogBBPolar surface area (Å²)*Plasma protein bindingPermeability glycoprotein

127543.52133.943950.918-3.27-1.96206.0767.321

* Calculated using Advanced Chemistry Development (ACD/Labs) Software (© 1994-2025 ACD/Labs)
Canonical SMILES22 other names for this Ligand

Canonical SMILES

O=C1C=CC=C(OC1C2C(=O)C=3C(O)=C4C(=C(O)C13)CC(O)(C(=O)CO)CC4OC5OC(C)C(O)C(N)C5

Isomeric SMILES

```
OC=C1C2=C([C@H](O)[C@H]3C[C@H](N)[C@H](O)[C@H](C)O3)C([C@H](C(CO)=O)
(O)C2)C(O)=C4C1C=O=C5C(C4=O)=C(OC)C=CC5
```

InChI
InChI=1S/C27H29NO11/c1-10-22(31)13(28)6-17(38-10)39-15-8-27(36,16(30)9-29)7-12-19(15)26(35)21-20(24(12)33)23(32)11-4-3-5-14(37-2)18(11)25(21)34/h3-5,10,13,15,17,22,29,31,33,35-36H,6-9,28H2,1-2H3/10-13,-15,-17,-22+27-m/s1

InChI Key
AOIJSUZBOXZONB-TZSSRYMLSA-N

22 other names for this Ligand

- (7S,9S)-7-[(2R,4S,5S,6S)-4-Amino-5-hydroxy-6-methyloxan-2-yl]oxy]-6,9,11-trihydroxy-9-(2-hydroxyacetyl)-4-methoxy-8,10-dihydro-7H-tetracene-5,12-dione
- (8S,10S)-10-[[3-Amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl]oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(2-hydroxyacetyl)-1-methoxy-5,12-naphthacenedione
- 14-Hydroxydaunomycin
- 5,12-Naphthacenedione, 10-[[3-amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl]oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-, (8S-cis)-
- 5,12-Naphthacenedione, 10-[[3-amino-2,3,6-trideoxy- α -L-lyxo-hexopyranosyl]oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-, (8S,10S)-

[View More](#) 

配體結構資訊與其他名稱

藥物情報資訊(Drug Intelligence)

CAS DRUG INTELLIGENCE

Approved Drug

Doxorubicin is an antineoplastic in the anthracycline class. General properties of drugs in this class include: interaction with DNA in a variety of different ways including intercalation (squeezing between the base pairs), DNA strand breakage and inhibition with the enzyme topoisomerase II. Most of...

View More

STATUS

Approval Status: US Approved
Approval Authority: US FDA
Approval Year: 1974, 2024
Originator: Farmitalia

INDICATIONS

Ovarian cancer
Modality: Primary
Highest Phase: Approved
Source

PRIMARY TARGET

DNA topoisomerase II
Function: Inhibitor
Measurement: IC50 2.67 µM
Source

Targets

View All

View Related Drugs

Primary Target

相關靶標

Function

Measurement

DNA topoisomerase II

Inhibitor

IC50 2.67 µM

Albumin

Binding Agent

-

DNA gyrase

Inhibitor

-

DNA topoisomerase II

Inhibitor

IC50 2.67 µM

A673

Inhibitor

IC50 200.0 nM

HepG2

Inhibitor

IC50 24.68 µM

Human ovarian cancer cell line

Inhibitor

IC50 0.074 µM

Indications

相關適應症

所處臨床階段和獲批資訊

X

Name	Modality	Highest Phase	Source
Ovarian cancer	Primary	Approved	View
AIDS-related Kaposi's sarcoma	Primary	Approved	View
Multiple myeloma	Primary	Approved	View
Locally advanced or unresectable soft tissue sarcoma	Primary	Phase III	View
Kaposi's sarcoma	Primary	Phase II	View
Glioblastoma	Primary	Phase II	View
Lung non-small cell carcinoma			View
Pancreatic ductal adenocarcinoma			View

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DOXIL, safely and effectively. See full prescribing information for DOXIL.

DOXIL® (doxorubicin HCl liposome injection) for intravenous infusion Initial U.S. Approval: 1995

WARNING: INFUSION REACTIONS, MYELOSUPPRESSION, CARDIOTOXICITY, LIVER IMPAIRMENT, SUBSTITUTION

See full prescribing information for complete boxed warning.

• Myocardial damage may lead to congestive heart failure and may occur as the total cumulative dose of doxorubicin HCl approaches 550 mg/m². Cardiac toxicity may also occur at lower cumulative doses with mediastinal irradiation or concurrent cardiotoxic agents (5.1).

• Acute infusion-related reactions, sometimes reversible upon terminating or slowing infusion, occurred in up to 10% of patients. Serious and sometimes fatal allergic/anaphylactoid like infusion reactions have been reported. Medication/emergency equipment to treat such reactions should be available for immediate use (5.2).

• Severe myelosuppression may occur (5.3).

• Reduce dosage in patients with impaired hepatic function (2.6).

• Accidental substitution of DOXIL resulted in severe side effects. Do not substitute on mg per mg basis with doxorubicin HCl (2.1).

RECENT MAJOR CHANGES

Contraindications, Nursing Mothers (4) Removed 8/2013

Warnings and Precautions, Secondary Oral Neoplasms (5.5) 8/2013

INDICATIONS AND USAGE

DOXIL is an anthracycline topoisomerase inhibitor indicated for:

• Ovarian cancer (1.1)

After failure of platinum-based chemotherapy.

• AIDS-related Kaposi's Sarcoma (1.2)

After failure of prior systemic chemotherapy or intolerance to such therapy.

• Multiple Myeloma (1.3)

In combination with bortezomib in patients who have not previously received bortezomib and have received at least one prior therapy.

DOSAGE AND ADMINISTRATION

Administer DOXIL as an initial rate of 1 mg/min to minimize the risk of infusion reactions. If no infusion related reactions occur, increase rate of infusion.

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: INFUSION REACTIONS, MYELOSUPPRESSION, CARDIOTOXICITY, LIVER IMPAIRMENT, ACCIDENTAL SUBSTITUTION

1 INDICATIONS AND USAGE

1.1 Ovarian Cancer

1.2 AIDS-Related Kaposi's Sarcoma

1.3 Multiple Myeloma

2 DOSAGE AND ADMINISTRATION

2.1 Usage and Administration Precautions

2.2 Patients With Ovarian Cancer

2.3 Patients With AIDS-Related Kaposi's Sarcoma

2.4 Patients With Multiple Myeloma

6 ADVERSE REACTIONS

6.1 Overall Adverse Reactions Profile

6.2 Adverse Reactions in Clinical Trials

6.3 Post Marketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

This label may not be the latest approved by FDA.

For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Infusion to complete administration over 1 hour. Do not administer as bolus injection or undiluted solution (2.1).

• Ovarian cancer: 50 mg/m² IV every 4 weeks for 4 courses minimum (2.2)

• AIDS-related Kaposi's Sarcoma: 20 mg/m² IV every 3 weeks (2.3)

• Multiple Myeloma: 30 mg/m² IV on day 4 following bortezomib which is administered at 1.3 mg/m² bolus on days 1, 4, 8 and 11, every 3 weeks (2.4)

DOSAGE FORMS AND STRENGTHS

Single use vial: 20 mg/10 mL and 50 mg/25mL (3)

CONTRAINDICATIONS

• Hypersensitivity reactions to a conventional formulation of doxorubicin HCl or the components of DOXIL (4, 5.3)

WARNINGS AND PRECAUTIONS

• Hand-Foot Syndrome may occur. Dose modification or discontinuation may be required (5.4)

• Radiation recall reaction may occur (5.5)

ADVERSE REACTIONS

Most common adverse reactions (≥20%) are asthenia, fatigue, fever, anorexia, nausea, vomiting, stomatitis, diarrhea, constipation, hand and foot syndrome, rash, neutropenia, thrombocytopenia and anemia (6).

To report SUSPECTED ADVERSE REACTIONS contact Janssen Products, LP at 1-800-JANSSEN (1-800-326-7736) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

• DOXIL may formulations

• DOXIL can

• Discontinue

See 17 for PATIENT COUNSELING INFORMATION.


Revised: 08/2013

15

© 2025 American Chemical Society. All rights reserved.

CAS
A division of the
American Chemical Society

配體藥理學資訊



← Prev (1 of 9,000) Next →

Target	Fibulin-1 (Protein)
Gene	FBLN1
Assay Type	Functional
Procedure	Inhibitory concentration of the compound against human MCF-7 cells transfected with a pool of two siRNA oligonucleotides specific for human fibulin-1
Function	Inhibitor
Parameter	IC50
pValue	13.05
Value	$9 \times 10^{-8} \mu\text{M}$
Experiment Type	in vitro
Organism	Homo sapiens
Cell	MCF-7
Source	Cancer Research (2007). 67(9). 4271-4277

實驗詳情

藥理活性實驗數據篩選項

Filters

Chemical Filters	^
▼ pValue	
Biological Filters	^
▼ Target	靶点
▼ Target Type	靶点类型
▼ Disease	疾病
▼ Organism	生物有機體
▼ Parameter	藥理學參數
▼ Function	功能
▼ Experiment Type	實驗類型
Source Filters	^
▼ Document Type	
▼ Publication Year	
▼ Language	

Organism

- ☐ Homo sapiens
- ☐ Mus musculus
- ☐ H. EP.2 ATCC CCL23
- ☐ Rattus norvegicus
- ☐ Sus scrofa

[View All](#)

Apply

Parameter

- ☐ IC50
- ☐ GI50
- ☐ Cell viability
- ☐ Inhibition
- ☐ Protein expression

[View All](#)

Apply

Function

- ☐ Inhibitor
- ☐ Modulator
- ☐ Binder
- ☐ Substrate
- ☐ Antagonist

[View All](#)

Apply

配體毒理學資訊

SummaryPharmacologyADMEToxicologyProteins (313)Diseases (301)Related Immunotherapeutics (12)Chemical SpaceMetabolites (4)Similar Ligands (764)

Applied Filters (0)

Parameter	Value	Organism	Assay	Source
ED (effective dose)	3.16228 × 10 ⁴ μM	Mus musculus	View	Journal of Medicinal Chemistry (2003), 46(14), 2985-3007
IC10	0.2 μM	Mus musculus	View	TIVIEQ Toxicology In Vitro. (Pergamon Press Inc., Maxwell House, Fairview Park, Elmsford, NY 10523) V.1- 1987- Volume(issue)/page/year
IC10	0.012 μM	Homo sapiens	View	TXAPA9 Toxicology and Applied Pharmacology. (Academic Press, Inc., 1 E. First St., Duluth, MN 55802) V.1- 1959- Volume(issue)/page/year
IC10	0.06 μM	Mus musculus	View	TIVIEQ Toxicology In Vitro. (Pergamon Press Inc., Maxwell House, Fairview Park, Elmsford, NY 10523) V.1- 1987- Volume(issue)/page/year
IC20	0.022 μM	Homo sapiens	View	TXAPA9 Toxicology and Applied Pharmacology. (Academic Press, Inc., 1 E. First St., Duluth, MN 55802) V.1- 1959- Volume(issue)/page/year
IC50	0.08 μM	Homo sapiens	View	AMCLCT ACS Medicinal Chemistry Letters. (American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036) V.1- 2010- Vo
IC50	0.48 μM	Homo sapiens	View	AMCLCT ACS Medicinal Chemistry Letters. (American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036) V.1- 2010- Vo
IC50	0.14 μM	Homo sapiens	View	TIVIEQ Toxicology In Vitro. (Pergamon Press Inc., Maxwell House, Fairview Park, Elmsford, NY 10523) V.1- 1987- Volume(issue)/page/year
IC50	1 mg/L	Homo sapiens	View	FRM 100 Pavia, Italy) V.44- 1989- Volume(issue)/page/year
IC50	4.8 mg/L	Mus musculus	View	JOET 100 Pavia, Ireland) V.1- 1979- Volume(issue)/page/year

← Prev12345...80Next →

Show10per page

Assay TypeFunctional

ProcedureLog cell kill of tumor bearing mice was determined by the dose of 162 mg/K

ParameterED (effective dose)

Parameter DetailLog Kill

pValue1.50

Value3.16228 × 10⁴ μM

Disease[Cancer](#)

Experiment Typein vivo

OrganismMus musculus

Source[Journal of Medicinal Chemistry \(2003\), 46\(14\), 2985-3007](#)

Biological Filters

毒理學實驗數據篩選項

Organism

☐ Homo sapiens

☐ Mus musculus

☐ Rattus norvegicus

☐ Mus

☐ Artemia salina

View All

生物有機體

Parameter

☐ LC50

☐ IC50

☐ LD50

☐ MTD

☐ LDLo

View All

毒理學參數

Experiment Type

☐ in vitro

☐ in vivo

Apply

實驗類型

Route of Administration

☐ Intraperitoneal

☐ Intravenous

☐ Oral

☐ Immersion

☐ Subcutaneous

Apply

給藥方式

Source Filters

Document Type

Publication Year

Language

18

© 2025 American Chemical Society. All rights reserved.

CAS

A division of the American Chemical Society

基於CAS數據關聯，一鍵關聯到該配體相關靶點與疾病資訊詳情頁，並探索更多關聯資訊

19

配體相關免疫療法資訊

Summary

Pharmacology

ADME

Toxicology

Proteins (313)

Diseases (301)

Related Immunotherapeutics (12)


Chemical Space

Metabolites (4)

Similar Ligands (764)

MAB-DMBA-SIL-Dox






Absolute stereochemistry shown

抗體與配體結構資訊

Antibody	Cetuximab
Antigen	Epidermal growth factor receptor (EGFR)
Payload	Doxorubicin
Therapeutic Target	DNA topoisomerase 2-alpha
Drug-Antibody Ratio	5.57
Linker	DMBA-SIL
Method	Site-specific conjugation with the respective EGFR mAb via thiolmaleimide Michael addition.

Trastuzumab-PC4AP-DOX





Absolute stereochemistry shown

Trastuzumab-PC4AP-DOX

Antibody	Trastuzumab
Antigen	Receptor tyrosine-protein kinase erbB-2 (ERBB2)
Payload	Doxorubicin
Therapeutic Target	DNA topoisomerase 2-alpha
Drug-Antibody Ratio	2
Linker	Photocaged C4AP
Method	Random conjugation through reduced inter-chain cysteines.

免疫療法詳情

免疫療法詳情

ADC藥物詳情

Trastuzumab emtansine

CAS Registry Number: 1018448-65-1

1,972


8

Retrosynthesis


Predictive Analytics

...

Summary Pharmacology ADME Proteins (2) Diseases (1)



抗體3D結構圖



藥物分子結構

Antibody	Trastuzumab
Antigen	Receptor tyrosine-protein kinase erbB-2 (ERBB2)
Payload	Mertansine DM1
Therapeutic Target	Microtubule
Drug-Antibody Ratio	3.5
Linker	Succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC)
Method	Random conjugation through nucleophilic lysines.

ADC藥物詳情

CAS DRUG INTELLIGENCE

藥物情報資訊

Approved Drug

Trastuzumab emtansine (ado-trastuzumab emtansine, trade name Kadcyla) is a combination between a monoclonal antibody and a small-molecule drug. Each molecule of trastuzumab emtansine consists of a single trastuzumab molecule with several molecules of DM1, a cytotoxic maytansinoid, attached. SMCC, or...

[View More](#)

STATUS	Approval Status: US Approved Approval Authority: US FDA Approval Year: 2013 Originator: Genentech
INDICATIONS	Breast cancer Modality: Primary Highest Phase: Approved Source
PRIMARY TARGET	Receptor tyrosine-protein kinase erbB-2 Function: Blocker Source

Heavy Chain Sequence

EVQLVESGGGLVQPGGSLRLSCAASG**GFNIKDTY**IHWVRQAPGKGLEWVAR**IYPTNGYT**TRYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYY**SRWGGDG**
FYAMDYDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL
TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKL
TVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Light Chain Sequence

DIQMTQSPSSLS**SAS**VGDRTVITCRAS**QDVNTA**AVAWYQQKPGKAPKLLIYSASFLYSGVPSRFSGSRSGETDFTLTISLQPEDFATYY**QQHYTTPPT**PTFGQG
TKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSYSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPV
TKSFNRGEC

重鏈/輕鏈序列

HCDR1
GFNIKDTY

HCDR2
IYPTNGYT

HCDR3
SRWGGDGFYAMDY


LCDR2
SAS

LCDR1
QDVNTA

LCDR3
QQHYTTPPT

CDR序列

相似配體結構與可視化分析



探索配體代謝產物

已知代謝產物：Known

預測代謝產物：Confidence Score置信度分值，
表示發生該生物轉化的概率

Doxorubicin
CAS Registry Number: 23214-92-8

112K 1,642 Retrosynthesis Predictive Analytics ...

Summary Pharmacology ADME Toxicology Proteins (313) Diseases (301) Related Immunotherapeutics (12) Chemical Space **Metabolites (4)** Similar Ligands (764)

24385-10-2 Known

54193-28-1 Known

69417-09-0 Known

Known

Transformation of 23214-92-8 to 24385-10-2

查看生物轉化反應

Facilitator	Organism	Source
-	Homo sapiens	Chemical Research in Toxicology (2000), 13(5), 414-420

大綱

- CAS BioFinder簡介與檢界面介紹（介紹與介面）
- 配體檢索與配體詳情信息（配體與細節概述）
- **骨架檢索與構效關係分析（支架與搜救分析）**
- 蛋白/通路/靶點、疾病與生物標誌物（Proteins and Pathways; 疾病與生物標記）
- 生物活性預測（Predictive Analytics）


基於靶點或結構片段，探索活性結構骨架

[Ligands](#)
[Scaffolds](#)
[Proteins](#)
[Diseases](#)

Search by disease name, ligand, protein, or draw a structure.

繪製核心結構片段

Draw a Structure



Filters

Chemical Filters

Rule of 5 Filter Preset ☐ OFF This will automatically be applied.

^ pValue
1.70 to 13.00

Reset Filter

Modality

Druglikeness

ADME

Commercial Availability

Approval Status

Approval Authority

Biological Filters

Target

☐ Menin

☐ Cyclin E1-CDK2 complex

☐ Histone-lysine N-methyltransferase complex

☐ GTPase KRas

☐ Monoglyceride lipase

[View All](#)

Scaffolds search for drawn structure

Ligands Scaffolds Pharmacology

2,649 Results

1 Ligands: 210 Proteins: 9

2 Ligands: 11 Proteins: 5

3 Ligands: 55 Proteins: 1

4 Ligands: 20 Proteins: 4

5 Ligands: 34 Proteins: 1

6 Ligands: 12 Proteins: 3

7 Ligands: 8 Proteins: 12

8 Ligands: 31 Proteins: 1

10 Ligands: 1 Proteins: 1

11 Ligands: 1 Proteins: 1

12 Ligands: 1 Proteins: 1

歸一化的活性

篩選關注的靶點

$$pKa = -\log_{10} Ka$$

$$pIC50 = -\log_{10} IC50$$

IC50越小，
pValue越大，抑
制活性越高。

探索目標結構骨架的SAR數據



Scaffolds


Search by protein, ligand, disease or draw a structure.



Edit

[Return to Results](#)

Scaffold Summary



Associated Proteins

1

▼

Filters

Chemical Filters

▲

OFF

Rule of 5 Filter Preset

This will automatically be applied.

▼

R-Groups

▼

pValue

▼

Druglikeness

▼

ADME

▼

Commercial Availability

Biological Filters

▲

▼

Target

▼

Target Type

▼

Disease



GIELXZVQJYHYPI-UHFFFAOYSA-N
Murcko scaffold InChI Key

Ligands

Pharmacology


Similar Scaffolds

☐

60 Results

☐ 1

[2765314-82-5](#)
[7-(2,5-Difluoro-3-methoxyphenyl)-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3-methy...




[Metabolites: 4](#)

[Proteins: 1](#)

☐ 2

[2765312-35-2](#)
[7-[3-Ethoxy-5-(trifluoromethyl)phenyl]-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3...




[Metabolites: 4](#)

[Proteins: 1](#)

☐ 3

[2765312-62-5](#)
[7-(3,5-Dimethylphenyl)-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3-methylcyclobuty...




[Metabolites: 4](#)

[Proteins: 1](#)

☐ 4

[2765312-53-4](#)
(cis-3-Hydroxy-3-methylcyclobutyl)[7-(2-methylphenyl)-2-azaspiro[3.5]non-2-yl]me...



[Metabolites: 7](#)

[Proteins: 1](#)

☐ 5

[2765312-61-4](#)
[7-[3-Ethyl-5-(trifluoromethyl)phenyl]-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3-methylcyclobutyl][7-(4-methylphenyl)-2-azaspiro[3.5]non-2-yl]me...




[Metabolites: 6](#)

[Proteins: 1](#)

☐ 7

[2765314-60-9](#)
[7-(3,5-Dimethylphenyl)-7-methoxy-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3-methy...




[Metabolites: 6](#)

[Proteins: 1](#)

☐ 8

[2765314-68-7](#)
[7-(2-Fluoro-3-methylphenyl)-2-azaspiro[3.5]non-2-yl][cis-3-hydroxy-3-methylcyclobutyl]me...



[Metabolites: 5](#)

[Proteins: 1](#)

Ligands: 60

Proteins: 1

Sort: Relevance ▼

 Get MMPA

R-Groups

R1

7

R2


13

R3

8

R4

探索目標結構骨架的SAR數據



符合分子對分析 (MMPA)

← [Return to Results](#)

Scaffold Summary

GIELXZVQJYHYPI-UHFFFAOYSA-N
Murcko scaffold InChI Key

Ligands Pharmacology Similar Scaffolds

☐ 60 Results

Sort: Relevance [Get MMPA](#)

☐ 1 **2765314-82-5**
[7-(2,5-Difluoro-3-methoxyphenyl)-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methy...

☐ 2 **2765312-35-2**
[7-[3-Ethoxy-5-(trifluoromethyl)phenyl]-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methy...

☐ 3 **2765312-62-5**
[7-(3,5-Dimethylphenyl)-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methylcyclobuty...

☐ 4 **2765312-53-4**
(cis-3-Hydroxy-3-methylcyclobutyl)[7-(2-methylphenyl)-2-azaspiro[3.5]non-2-yl]me...

Associated Proteins
1

Filters

Chemical Filters

☐ Rule of 5 Filter
This will automati...

Select a Target for Matched Molecular Pairs Analysis (MMPA)

☐ **Monoglyceride lipase**

選擇目標靶點

Select one R-Group Cluster to Visualize

15 Matched Molecular Pairs

10 Matched Molecular Pairs

10 Matched Molecular Pairs

10 Matched Molecular Pairs

獲取目標R基團位點變化的MMPA活性對比結果


Show 100 per page

[Previous](#) [Submit](#) [Cancel](#)

[Next](#) [Cancel](#)

符合分子對分析 (MMPA)


快速直觀獲取匹配分子對 $\Delta pActivity$ 熱圖



查看同一結構骨架中的配體藥理活性

[Return to Results](#)

Scaffold Summary



Associated Proteins
1

Filters

Chemical Filters

Biological Filters

Target

☒ Monoglyceride lipase

Target Type

Disease

Organism

Parameter

Experiment Type

Source Filters

Document Type

Publication Year

GIELXZVQJYHYPI-UHFFFAOYSA-N

Murcko scaffold InChI Key

Ligands




Pharmacology

Similar Scaffolds




Applied Filters (1) Target: Monoglyceride lipase X Clear all filters

Show Structures ON

Ligand	Target	Gene	Parameter	pValue	Value	Function	Assay	Source
 2765314-60-9 [7-(3,5-Dimethylphenyl)-7-methoxy-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methy...	Monoglyceride lipase (Protein)	MGLL	IC50	9.08	$8.4 \times 10^{-4} \mu\text{M}$	-	View	United States, US20220089538 A1 2022-03-24
 2765313-81-1 (cis-3-Hydroxy-3-methylcyclobutyl)[7-methoxy-7-[3-(1-methylethyl)phenyl]-2-azasp...	Monoglyceride lipase (Protein)	MGLL	IC50	8.92	0.0012 μM	-	View	United States, US20220089538 A1 2022-03-24
 2765314-75-6 (cis-3-Hydroxy-3-methylcyclobutyl)[7-[2-methyl-3-(1-methylethoxy)phenyl]-2-azasp...	Monoglyceride lipase (Protein)	MGLL	IC50	8.82	0.0015 μM	-	View	United States, US20220089538 A1 2022-03-24

Assay Detail




Ligand
[2765314-60-9](#)


C23H33NO3
[7-(3,5-Dimethylphenyl)-7-methoxy-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methy...

實驗詳情

Target	Monoglyceride lipase (Protein)
Gene	MGLL
Assay Type	Functional
Procedure	FAAH assay
Parameter	IC50
pValue	9.08
Value	$8.4 \times 10^{-4} \mu\text{M}$
Disease	Disease of cellular proliferation
Experiment Type	in vitro
Organism	Homo sapiens
Cell	HeLa
Cell Detail	Cervical cancer cell
Source	United States, US20220089538 A1 2022-03-24

藥理活性數據


Ligand	Target	Gene	Parameter	pValue	Value	Function	Assay	Source
 2765314-60-9 [7-(3,5-Dimethylphenyl)-7-methoxy-2-azaspiro[3.5]non-2-yl](cis-3-hydroxy-3-methy...	Monoglyceride lipase (Protein)	MGLL	IC50	9.08	$8.4 \times 10^{-4} \mu\text{M}$	-	View	United States, US20220089538 A1 2022-03-24
 2765313-81-1 (cis-3-Hydroxy-3-methylcyclobutyl)[7-methoxy-7-[3-(1-methylethyl)phenyl]-2-azasp...	Monoglyceride lipase (Protein)	MGLL	IC50	8.92	0.0012 μM	-	View	United States, US20220089538 A1 2022-03-24
 2765314-75-6 (cis-3-Hydroxy-3-methylcyclobutyl)[7-[2-methyl-3-(1-methylethoxy)phenyl]-2-azasp...	Monoglyceride lipase (Protein)	MGLL	IC50	8.82	0.0015 μM	-	View	United States, US20220089538 A1 2022-03-24

30

© 2025 American Chemical Society. All rights reserved.









CAS
A division of the
American Chemical Society

獲取與目標骨架相似骨架結構，啟發新的SAR假設

[Return to Results](#)
Scaffold Summary

Associated Proteins
1

GIELXZVQJYHYPI-UHFFFAOYSA-N
Murcko scaffold InChI Key
Ligands Pharmacology Similar Scaffolds
1,123 Scaffolds

**獲取相似骨架結構(Similar Scaffolds) ;
基於相應藥效團的關鍵結構元素識別相似骨架結構**

1  Ligands: 2 Proteins: 1	2  Ligands: 9 Proteins: 2	3  Ligands: 3 Proteins: 1	4  Ligands: 9 Proteins: 2
5  Ligands: 69 Proteins: 1	6  Ligands: 1 Proteins: 1	7  Ligands: 9 Proteins: 1	8  Ligands: 2 Proteins: 1

大綱

- CAS BioFinder簡介與檢界面介紹（介紹與介面）
- 配體檢索與配體詳情信息（配體與細節概述）
- 骨架檢索與構效關係分析（支架與搜救分析）
- 蛋白/通路/靶點、疾病與生物標誌物（**Proteins and Pathways**; 疾病與生物標記）
- 生物活性預測（Predictive Analytics）

通過靶點、疾病或配體，獲取歸一化的蛋白檢索結果

CAS BioFinder

Proteins SYK 示例：SYK

Filters

- Chemical Filters
- Biological Filters
 - Protein
 - Disease
 - Organism
 - Parameter
 - Function
 - Experiment Type
- Source Filters
 - Document Type
 - Publication Year
 - Language

Proteins search for SYK

Proteins Pharmacology

4 Results

Sort: Relevance View: **Grid** (Grid, List)

1	2	3	4
SYK (Homo sapiens) Tyrosine-protein kinase SYK	SYK Tyrosine-protein kinase SYK	Syk (Rattus norvegicus) Tyrosine-protein kinase SYK	Syk (Mus musculus) Tyrosine-protein kinase SYK
Diseases: 223 Ligands: 17K	Diseases: 344 Ligands: 16K	Diseases: 15 Ligands: 546	Diseases: 13 Ligands: 35

Return to top

切換縮圖檢視或清單檢視


蛋白詳情資訊與立體結構展示

SYK

Tyrosine-protein kinase SYK

Summary Pharmacology Ligands (31K) Diseases (388) Chemical Space Pathways

Crystal structure of the syk tyrosine kinase domain with Staurosporin



蛋白的三維晶體結構及與其相互作用的小分子

STAUROSPORINE

1XBC | Model 1 | Instance ASML1 | B [auth A] | STU 1

1XBC

↺

+

-

↓

下載cif格式結構檔

Homo sapiens

unspecified

Rattus norvegicus

Mus musculus

來源有機體

Organism

Homo sapiens

GET FASTA

Copy Sequence

>P43405-1|Tyrosine-protein kinase SYK|Homo sapiens

MASSGMADSANHLPPFFGNITREEAEDYLQGGMSDGLYLLRQSRNYLGGFALSVAHGRKAHHYIERELNGTYAIAGGRTHASPADLCHYHSQESDGLVCLLKPKFNRPGVQVQPK
TGPFEDLKENLIREYVKQTWNLQGALEQAIISQKPQLEKIATTAHEKMPWFHGKISREESEQIVLIGSKTNGKFLIRARDNNGSYALCLLHEGKVLHYRIDKDTGKLSIPEGK
KFDTLWQLVEHYSYKADGLLRVLTPCQKIGTQGNVNFGRPQLPGSHPATWSAGGISRIKSYSFPPKPGHRKSSPAQGNRQESTVSFNPYEPELAPWAADKGPQREALPMDTEVY
ESPYADPEEIRPKEVYLDRKLLTLEDKELGSGNFGTVKKGYQMKVVKTVAVKILKNEANDPALDELLAEANVMQQLDNPIYIRMIIGICEASWMLVMEMAEGLPLNKYLQQR
HVKDKNIIELVHQVSMGMKYLEESNFVHRDLAARNVLLVTQHYAKISDFGLSKALRADENYYKAQTHGKWPVKWYAPECINYYKFSSKSDVWSFGVLMWEAFSYGQKPYRGMKGSE
VTAMLEKGERMGCPCAGCPREMYDLMLNCWTYDVENRPGFAAVELRLRNYYDVVN

序列詳情

Identifier	Method	Resolution	Chain	Positions	Source
1A81	X-Ray	3.0 Å	A/C/E/G/I/K	9 - 262	PDB
1CSY	NMR	-	A	163 - 265	PDB
1CSZ	NMR	-	A	163 - 265	PDB
1XBA	X-Ray	2.00 Å	A	353 - 635	PDB
1XBB	X-Ray	1.57 Å	A	353 - 635	PDB
1XBC	X-Ray	2.00 Å	A	353 - 635	PDB
3BUW	X-Ray	1.450 Å	A/C	317 - 329	PDB
				349 - 635	PDB

External Links and Identifiers

Alliance of Genome Resources: [HGNC:11491](#)

ChEMBL: [CHEMBL2599](#)

Ensembl gene: [ENSG00000165025.15](#), [ENSG00000165025](#)

HGNC: [HGNC:11491](#)

NCBI Gene: [6850](#)

OMIM: [600085](#)

Uniprot reviewed: [P43405](#)

[View Less](#)

歸一化的靶點名稱
與其他詳情資訊

Synonyms

Spleen associated tyrosine kinase
Spleen tyrosine kinase

相關配體與藥理活性數據



相關配體

Summary Pharmacology **Ligands (31K)** Diseases (388) Chemical Space Pathways

Sort: Relevance

1	841290-80-0 6-[[[5-Fluoro-2-[(3,4,5-trimethoxyphenyl)amino]-4-pyrimidinyl]amino]-2,2-dimethyl...	Metabolites: 2	Proteins: 581
2	62996-74-1 Staurosporine	Metabolites: 4	Proteins: 1,024
3	1370261-96-3 2-[[[(1R,2S)-2-Aminocyclohexyl]amino]-4-[[3-(2H-1,2,3-triazol-2-yl)phenyl]amino]-...	Metabolites: 0	Proteins: 27
4	152459-95-5 Imatinib	Metabolites: 1	Proteins: 676
5	544444-90-8 (-)-cis-Tetrahydroisohumulone		
6	732983-37-8 2-[[[7-(3,4-Dimethoxyphenyl)imidazo[1,2-c]pyrimidin-5-yl]amino]-3-pyridinecarboxy...		
7	1240390-27-5 2-[[[(3R,4R)-3-Aminotetrahydro-2H-pyran-4-yl]amino]-4-[[4-methylphenyl]amino]-5-p...		
8	841290-81-1 Tamatinib		

Ligand	Parameter	pvalue	value	function	Organism	Assay	Source
 1438423-22-3 5-[[[(1R,2S)-2-Aminocyclohexyl]amino]-3-(3-pyridinylamino)-2-pyrazinecarboxamide	IC50	12.00	1 × 10 ⁻⁶ μM	Inhibitor	Homo sapiens	View	United States, US20130131040 A1 2013-05-23
 1438421-77-2 5-[[[(1R)-1-(Aminocarbonyl)propyl]amino]-3-(2,1-benzisothiazol-3-ylamino)-2-pyrazinecarboxamide	IC50	12.00	1 × 10 ⁻⁶ μM	Inhibitor	Homo sapiens	View	United States, US20130131040 A1 2013-05-23
 1438422-34-4 5-[[[(1R,2S)-2-Aminocyclohexyl]amino]-3-[[3-methyl-5-isothiazolyl]amino]-2-pyrazinecarboxamide	IC50	12.00	1 × 10 ⁻⁶ μM	Inhibitor	Homo sapiens	View	United States, US20130131040 A1 2013-05-23
 1438420-30-4 5-[[[(1R)-1-(Aminocarbonyl)-3,3-difluoropropyl]amino]-3-[[3,4-dimethylphenyl]amino]-2-pyrazinecarboxamide	IC50	12.00	1 × 10 ⁻⁶ μM	Inhibitor	Homo sapiens	View	United States, US20130131040 A1 2013-05-23

蛋白相關疾病資訊




SYK
Tyrosine-protein kinase SYK

Summary Pharmacology Ligands (31K) **Diseases (388)** Chemical Space Pathways

Sort: Relevance ▾ [View all associated diseases](#)

Disease	疾病	Description	疾病描述	Biomarkers	生物標誌物
1	Cancer	A disease of cellular proliferation that is malignant and primary, characterized by uncontrolled cellular proliferation, local cell invasion and metastasis.		BRCA1 DNA repair associated, Neurotrophin 4, Autophagy related 16 like 1, Major histocompatibility complex, class II, DP beta 1, KRAS proto-oncogene, GTPase, View All	7,479 1M
2	Alzheimer's disease	A tauopathy that is characterized by memory lapses, confusion, emotional instability and progressive loss of mental ability and results in progressive memory loss, impaired thinking, disorientation, and changes in personality and mood starting and leads in advanced cases to a profound decline in cognitive and physical functioning and is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid.		Apolipoprotein E, Fibrinogen alpha chain, Islet amyloid polypeptide, APP (rs199862130) polymorphism, Synuclein alpha, View All	4,974 722K
3	Rheumatoid arthritis	An arthritis that is an autoimmune disease which attacks healthy cells and tissue located in joint.		TIMP metalloproteinase inhibitor 1, Solute carrier family 22 member 4, Tumor necrosis factor, Matrix metalloproteinase 1, Protein kinase C theta, View All	3,847 576K
4	Asthma	A bronchial disease that is characterized by chronic inflammation and narrowing of the airways, which is caused by a combination of environmental and genetic factors. The disease has symptom recurring periods of wheezing (a whistling sound while breathing), has symptom chest tightness, has symptom shortness of breath, has symptom mucus production and has symptom coughing.		Membrane spanning 4-domains A2, Transient receptor potential cation channel subfamily A member 1, Adrenomedullin, Fc epsilon receptor 1a, CDKN2B-AS1 (rs7859362) polymorphism, View All	3,333 516K
5	Diabetes mellitus	A glucose metabolism disease that is characterized by chronic hyperglycaemia with disturbances of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action, or both.		MTHFR (rs1801133) polymorphism, Cytokine inducible SH2 containing protein, Cystatin C, Insulin, Mechanistic target of rapamycin kinase, View All	3,871 401K

蛋白的生物通路資訊



SYK
Tyrosine-protein kinase SYK

[Summary](#) [Pharmacology](#) [Ligands \(31K\)](#) [Diseases \(388\)](#) [Chemical Space](#) **[Pathways](#)**

Pathway	Reaction	Source
alpha IIb beta3 pathway	Fibrinogen:(alpha IIb beta3)2:(Src)2 + 2 CIB:Ca + 2 talin <=> Fibrinogen:(alpha IIb beta3)2:(Src)2:(CIB)2:(Ca)2:(talin)2	Proceedings of the National Academy of Sciences of the United States of America (2003), 100(23), 13298-13302 Journal of Biological Chemistry (1997), 272(8), 4651-4654
	Rac1:GDP + GTP --VAV1{pY}:SLP-76{p}:Nck-1--> Rac1:GTP + GDP	EMBO Journal (1998), 17(22), 6608-6621 Journal of Biological Chemistry (2001), 276(8), 5916-5923
	alpha IIb beta3 + Csk + Src <=> alpha IIb beta3:Csk:Src	Journal of Cell Biology (2002), 157(2), 265-275
	Fibrinogen:(alpha IIb beta3)2:(Src{pY419})2:(CIB)2:(Ca)2:(talin)2:(Syk{pY})2 + 2 NDP	Coordinate interactions of Csk, Src, and Syk kinases with alphaIIb beta3 initiate integrin signaling to the cytoskeleton Biology (2002), 157(2), 265-275
	Fibrinogen + 2 alpha IIb beta3:Csk:Src{pY529}	Biology (2002), 157(2), 265-275
	SLP-76 + NTP --Fibrinogen:(alpha IIb beta3)2:	Journal of Biological Chemistry (2001), 276(8), 5916-5923 Journal of Biological Chemistry (1996), 271(42), 25746-25749
	vasp + SLAP-130{p} + SLP-76{p} <=> vasp:SLP-76{p}	Journal of Biological Chemistry (1995), 270(49), 29071-4 Journal of Biological Chemistry (2001), 276(8), 5916-5923
	Fibrinogen:(alpha IIb beta3)2:(Src{pY529})2 --	
	VAV1{pY}:SLP-76{p}:Nck-1 + PAK1 + Rac1:GTP	

Coordinate interactions of Csk, Src, and Syk kinases with alphaIIb beta3 initiate integrin signaling to the cytoskeleton

By: Oberfell, Achim; Eto, Koji; Mocsai, Attila; Buensuceso, Charito; Moores, Sheri L.; Brugge, Joan S.; Lowell, Clifford A.; Shattil, Sanford J.

DOI: 10.1083/jcb.200112113

Integrins regulate cell adhesion and motility through tyrosine kinases, but in contrast to the behavior of Src and Csk, Syk was associated with alphaIIb beta3 only after fibrinogen binding. Platelets multiply deficient in Src, Hck, Fgr, and Lyn, or normal platelets treated with Src kinase inhibitors failed to spread on fibrinogen. Inhibition of Src kinases blocked Syk activation and inhibited phosphorylation of Syk substrates (Vav1, Vav3, SLP-76) implicated in cytoskeletal regulation. Syk-deficient platelets exhibited Src activation upon adhesion to fibrinogen, but no spreading or phosphorylation of Vav1, Vav3, and SLP-76. These studies establish that platelet spreading on fibrinogen requires sequential activation of Src and Syk in proximity to alphaIIb beta3, thus providing a paradigm for initiation of integrin signaling to the actin cytoskeleton.

Keywords: Csk Src Syk kinase integrin signaling cytoskeleton platelet adhesion

[View Source](#) [Full Text](#)

Publication Information • Journal

Source	Database Information	Company/Organization	Publisher	Language
Journal of Cell Biology Volume: 157 Issue: 2 Pages: 265-275 Journal: Article: Research Support, Non-U.S. Govt; Research Support, U.S. Govt; P.H.S. 2002 DOI: 10.1083/jcb.200112113	AN: 2002:296764 CAN: 137:106866 PubMed ID: 11940607 CAPUS and MEDLINE	Division of Vascular Biology, Department of Cell Biology The Scripps Research Institute La Jolla, California 92037 United States	Rockefeller University Press	English

37

© 2025 American Chemical Society. All rights reserved.



CAS
A division of the
American Chemical Society

通過疾病名稱、靶點或配體檢索相關疾病資訊

CAS BioFinder

Diseases APP

×

Q

Draw

← Return to all Filters

Biomarker

Enter a Biomarker... Search

0 Selected

☐ Insulin

☐ Hemoglobin subunit alpha 2

☐ Hemoglobin subunit alpha 1

☐ Hemoglobin subunit beta

☐ Epidermal growth factor receptor

☐ C-reactive protein

☐ Erb-b2 receptor tyrosine kinase

☐ Microtubule-associated protein tau

☐ Estrogen receptor

☐ Tumor necrosis factor

☐ Interleukin 6

☐ Apolipoprotein E

☐ Kallikrein related peptidase 3

☐ Synuclein alpha

☐ KRAS proto-oncogene, GTPase

☐ Adiponectin, C1Q and collagen domain containing

☐ BRCA1 DNA repair associated

☐ AKT serine/threonine kinase 1

☐ Androgen receptor

☐ Leptin

☐ Vascular endothelial growth factor A

☐ Glucagon

☐ Angiotensin I converting enzyme

☐ Phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha

☐ Amyloid beta precursor protein

Diseases search for APP

Diseases

172 Results

疾病名稱

疾病簡介

生物標誌物

相關靶點與配體

Sort: Relevance

Disease	Description	Biomarkers		
1 Cancer	A disease of cellular proliferation that is malignant and primary, characterized by uncontrolled cellular proliferation, local cell invasion and metastasis.	BRCA1 DNA repair associated, Neurotrophin 4, Autophagy related 16 like 1, Major histocompatibility complex, class II, DP beta 1, KRAS proto-oncogene, GTPase, View All	7,479	1M
2 Alzheimer's disease	A tauopathy that is characterized by memory lapses, confusion, emotional instability and progressive loss of mental ability and results in progressive memory loss, impaired thinking, disorientation, and changes in personality and mood starting and leads in advanced cases to a profound decline in cognitive and physical functioning and is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid.	Apolipoprotein E, Fibrinogen alpha chain, Islet amyloid polypeptide, APP (rs199862130) polymorphism, Synuclein alpha, View All	4,974	722K
3 Rheumatoid arthritis	An arthritis that is an autoimmune disease which attacks healthy cells and tissue located in joint.	TIMP metalloproteinase inhibitor 1, Solute carrier family 22 member 4, Tumor necrosis factor, Matrix metalloproteinase 1, Protein kinase C theta, View All	3,847	576K
4 Asthma	A bronchial disease that is characterized by chronic inflammation and narrowing of the airways, which is caused by a combination of environmental and genetic factors. The disease has symptom recurring periods of wheezing (a whistling sound while breathing), has symptom chest tightness, has symptom shortness of breath, has symptom mucus production and has symptom coughing.	Membrane spanning 4-domains A2, Transient receptor potential cation channel subfamily A member 1, Adrenomedullin, Fc epsilon receptor 1a, CDKN2B-AS1 (rs7859362) polymorphism, View All	3,333	516K
5 Diabetes mellitus	A glucose metabolism disease that is characterized by chronic hyperglycaemia with disturbances of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action, or both.	MTHFR (rs1801133) polymorphism, Cytokine inducible SH2 containing protein, Cystatin C, Insulin, Mechanistic target of rapamycin kinase, View All	3,871	401K
6 Bacterial infectious disease	A disease by infectious agent that results in infection, has material basis in Bacteria.	Tumor necrosis factor, Activation induced cytidine deaminase, Serum amyloid A1, Transforming growth factor beta 1, C-C motif chemokine ligand 17, View All	2,645	202K
7 Parkinson's disease	A synucleinopathy that has material basis in degeneration of the central nervous system that often impairs motor skills, speech, and other functions.	Synuclein alpha, Leucine rich repeat kinase 2, Ubiquitin C-terminal hydrolase L1, Glucosylceramidase beta 1, Mechanistic target of rapamycin kinase, View All	4,497	423K
8 Psoriasis	A skin disease that is characterized by patches of thick red skin and silvery scales.	Interleukin 1 beta, Insulin, C-C motif chemokine ligand 5, Acyl-CoA thioesterase 12, ADAM metalloproteinase domain 10, View All	2,937	430K

疾病詳情頁與相關藥物資訊

Alzheimer's disease

相關生物標誌物、藥理活性數據、配體與靶點

Summary Biomarkers Pharmacology Ligands (705K) Proteins (4,937)

A tauopathy that is characterized by memory lapses, confusion, emotional instability and progressive loss of mental ability and results in progressive memory loss, impaired thinking, disorientation, and changes in personality and mood starting and leads in advanced cases to a profound decline in cognitive and physical functioning and is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid.

疾病簡介

Synonyms

Aging-related Alzheimer's disease
Alzheimer disease
Alzheimer's
Alzheimer's senile dementia
Alzheimer's type dementia
Alzheimers dementia
Early onset Alzheimer's disease
Intermediate-stage Alzheimer's disease
Late onset Alzheimer's disease

[View Less](#) ^

External Links and Identifiers

GARD: [10254](#) [↗](#)
KEGG: [05010](#) [↗](#)
MESH: [D000544](#) [↗](#)
NCI: [C2866](#) [↗](#)

歸一化的疾病
同義詞清單

AI Drug Intelligence Summarization

AI總結藥物情報資訊

Here is a summary of the state of drugs used to treat Alzheimer's disease:

CAS BioFinder shows 327 drug entries targeting Alzheimer's disease, predominantly consisting of small molecule compounds. The drugs exhibit diverse mechanisms of action, targeting various biological pathways including cholinergic signaling (acetylcholinesterase inhibitors, neuronal acetylcholine receptor agonists), kinase pathways (tyrosine kinase inhibitors targeting c-Kit, PDGFR, and Lyn), neurotransmitter systems (sigma opioid receptor agonists, serotonin 5-HT4 receptor agonists), and cellular signaling (glycogen synthase kinase-3 beta inhibitors, voltage-gated calcium channel blockers). The therapeutic approaches include both primary treatments aimed at disease modification and palliative treatments for symptom management. Development phases range from early Phase I studies to Phase III trials, with some compounds having achieved approved status. Target functions encompass inhibitors, agonists, partial agonists, and blockers, reflecting the multifaceted approach to addressing the complex pathophysiology of Alzheimer's disease through modulation of neuroinflammation, neurotransmission, and neuroprotective pathways.

This is an AI-generated summary of the drug intelligence information, and may be incomplete or include some inaccuracies. Please consult the records in the table for the source empirical data.

CAS BioFinder AI is always improving. [suggest improvements](#), or [learn more](#).

DRUGS

藥物情報資訊清單

Ligand ↕	Modality ↕	Highest Phase ↕ 1	Source
98185-20-7 Raclopride tartrate	Diagnostic	Approved	-
1098-97-1 Pyritinol	Primary	Approved	Dementia (Base, Switzerland), 0_5(2)_88-98
23173-12-8 <i>rel</i> -(4a <i>R</i> ,6 <i>S</i> ,8a <i>R</i>)-4a,5,9,10,11,12-Hexahydro-3-methoxy-11-methyl-6 <i>H</i> -benzofuro[3a,3...	Palliative	Approved	Source ↗
86332-23-2 1,2,3-Propanetriol, 2-(dihydrogen phosphate), monopotassium salt	Palliative	Approved	Source ↗
120011-70-3 Donepezil hydrochloride	Palliative	Approved	Source ↗

疾病與生物標誌物

✦ AI Table Summarization

There are 24 top biomarkers for Alzheimer's disease:
Select a biomarker to see more details.
[Microtubule associated protein tau](#), [Amyloid beta precursor protein](#), [Apolipoprotein E](#), [Tumor necrosis factor](#), [Insulin](#), [Brain derived neurotrophic factor](#), [Glycogen synthase kinase 3 beta](#), [Insulin like growth factor 1](#), [Triggering receptor expressed on myeloid cells 2](#), [Presenilin 1](#), [Acetylcholinesterase \(Yt blood group\)](#), [ATP binding cassette subfamily A member 7](#), [Butyrylcholinesterase](#), [Beta-secretase 1](#), [Sortilin related receptor 1](#), [Angiotensin I converting enzyme](#), [Granulin precursor](#), [Synuclein alpha](#), [Glutamate ionotropic receptor NMDA type subunit 2B](#), [Clusterin](#), [Interleukin 1 beta](#), [Presenilin 2](#), [Advanced glycosylation end-product specific receptor](#), [Translocase of outer mitochondrial membrane 40](#)

Alzheimer's disease

Summary **Biomarkers** Pharmacology Ligands (722K) Proteins (4,974)


Apply **生物標誌物** 參數 實驗說明 有機體 文獻來源

Show Filters Summarize

Biomarker	Gene	Category	Parameter	Measurement	Organism	Assay	Source
(-)-Thyroxine	-	Diagnostic	Biomarker level	Mean free Thyroxine (FT4) level was observed in 40 patients with alzheimer's disease with blunted TSH response which is significantly (P<0.01) more severely demented when compared to patients with non blunted TSH response	Human	View	Biological psychiatry (), 30(6), 567-76
(+)-Aldosterone	-	Pharmacodynamic/resp					Health Evaluation in African Americans Using RAS Therapy
(25R)-26-Hydroxycholesterol	-	Prognostic					Brain Communications (2023), 5(5), 1-14
[Pyruvate dehydrogenase (acetyl-transferring)] kinase isozyme 1, mitochondrial	-	Diagnostic					Journal of Biological Chemistry (2012), 287(44), 37245-37258

Biomarker

[51-48-9](#)



L-Tyrosine, O-(4-hydroxy-3,5-diiodophenyl)-3,5-diiodo-

Category

Diagnostic

Parameter

Biomarker level

Measurement

Mean free Thyroxine (FT4) level was observed in 40 patients with alzheimer's disease with blunted TSH response which is significantly (P<0.01) more severely demented when compared to patients with non blunted TSH response, Mean free Thyroxine(FT4) level was observed in 40 patients with alzheimer's disease having blunted TSH response which was significantly (P<0.03) higher when compared to patients with non blunted TSH response, High, Mean Thyroxine (T4) level was observed in 40 alzheimer's disease patients which is significantly (P<0.05) lower when compared to 17 elderly patients with major depression, Low

Organism

Human

Source

[Biological psychiatry \(\), 30\(6\), 567-76](#)

實驗詳情

40

© 2025 American Chemical Society. All rights reserved.

CAS
A division of the
American Chemical Society

大綱

- CAS BioFinder簡介與檢界面介紹（介紹與介面）
- 配體檢索與配體詳情信息（配體與細節概述）
- 骨架檢索與構效關係分析（支架與搜救分析）
- 蛋白/通路/靶點、疾病與生物標誌物（Proteins and Pathways; 疾病與生物標記）
- 生物活性預測（**Predictive Analytics**）

Predictive Analytics：通過演算法預測配體對不同靶點的生物活性數據

The image illustrates the workflow for creating a predictive analytics set in the CAS software. It starts with the main dashboard, which includes tabs for Ligands, Scaffolds, Proteins, and Diseases. A search bar is present, and a 'Predictive Analytics' section is highlighted with a yellow box, containing an icon and the text: 'Upload and analyze ligand data sets using algorithms to find bioactivity data insights.' An arrow points from this section to the 'Predictive Sets' panel.

The 'Predictive Sets' panel shows a 'History (405)' and 'Predictive Sets (24)' section. A 'Create New Set' button is highlighted with a yellow box. An arrow points from this button to the 'Create Predictive Analytics Set' dialog.

The 'Create Predictive Analytics Set' dialog has the following fields and options:

- Name your set:** Predictive_Set_09_09_2025_0952
- Select Color:** Light Blue (dropdown)
- Select Panel:** CAS Full Pharmacology (Default) (dropdown menu is open, showing options: CAS Full Pharmacology (Default), CAS Broad Safety Screen, CAS Medium Safety Screen, CAS Narrow Safety Screen, CAS Alzheimer's Disease Screen, CAS Breast Cancer Screen, CAS Colon Cancer Screen, CAS Depressive Disorder Screen)
- Upload a predefined set (Optional):** Select a file (.sdf) or drag and drop here. (A blue box labeled '上傳.sdf文件' points to this area)
- Buttons:** Create (highlighted with a yellow box), Cancel

A blue box labeled '選擇不同的預測面板' points to the 'Select Panel' dropdown menu.

An arrow points from the 'Predictive Test XXX' card in the 'Predictive Sets' panel to a context menu with the following options:

- Rerun analysis for set
- Add ligands from file
- Edit set
- Delete set

可從檢索結果集中添加配體結構

▼ Filters

Chemical Filters

OFF

Rule of 5 Filter Preset
This will automatically be applied.

▼ pValue

▼ Modality

▼ Druglikeness

▼ ADME

▼ Commercial Availability

▼ Approval Status

▼ Approval Authority

Biological Filters

▼ Target

▼ Target Type

▼ Disease

▼ Organism

▼ Parameter

▼ Function

▼ Experiment Type

Ligands search for drawn structure

Ligands Scaffolds Pharmacology

25 of 9,190 Selected

Sort: Relevance

Predictive Analytics

1

1961259-82-4

2,5-Diamino-1-(4-methylphenyl)-1H-pyrrole-3,4-dicarboxaldehyde



Metabolites: 2

Proteins: 0

2

1961259-76-6

2,5-Diazo-1-(4-methylphenyl)-1H-pyrrole-3,4-dicarboxaldehyde



Metabolites: 2

Proteins: 0

3

2270167-74-1

(3R,4R)-4-Methyl-(8-methyl-5-quinoliny)-3-pyrrolidinamine




Metabolites: 4

Proteins: 2

4

2002854-95-5

4-(3-Hydroxy-3-methyl-1-pyrrolidiny)-3-methylbenzonitrile



Metabolites: 4

Proteins: 2

5

2270167-41-2

8-(3-Amino-4-methyl-1-pyrrolidiny)-5-quinoxalinecarbonitrile




Metabolites: 4

Proteins: 2

6

2270167-40-1

8-[(3R,4R)-3-Amino-4-methyl-1-pyrrolidiny]-5-quinoxalinecarbonitrile




Metabolites: 4

Proteins: 2

7

1067895-50-4

8-Cyclobutyl-2-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decane



Metabolites: 4

Proteins: 2

×

Predictive Analytics

Create New Set Add to Existing Set

Name your set

Predictive_Set_09_09_2025_1002

Select Color

Light Blue

Select Panel

CAS Breast Cancer Screen

25 selected ligands will be added to this set

Create Cancel

43

© 2025 American Chemical Society. All rights reserved.

CAS
A division of the
American Chemical Society

不同的蛋白靶点面板说明

History (407) Predictive Sets (25)

Create New Set **Panels**

Predictive Set **Panels**

CAS Breast Cancer Screen

Ligands: 25

Updated on 09 September

+ Create New Panel

Search panels by name...

Panel Name	Description	Targets
CAS Full Pharmacology (Default)	The full panel of protein target models available through CAS to highlight potential safety concerns and treatment options	All
CAS Broad Safety Screen	A diverse panel of protein targets to widely test the safety of your drug candidate	192
CAS Medium Safety Screen	Panel of protein targets likely to be implicated to adverse drug reactions (ADRs) based off of known structure activity relationships (SARs)	115
CAS Narrow Safety Screen	Panel of protein targets established as being involved in adverse drug reactions (ADRs). These targets are essential for early safety screening of drugs	49
CAS Alzheimer's Disease Screen	Panel of protein targets potentially implicated in Alzheimer's Disease	81
CAS Breast Cancer Screen	Panel of protein targets potentially implicated in breast cancer and neoplasms	70
CAS Colon Cancer Screen	Panel of protein targets potentially implicated in colon cancer and neoplasms	92
CAS Depressive Disorder Screen	Panel of protein targets potentially implicated in depressive disorder and major depressive disorder	51
CAS Diabetes Screen	Panel of protein targets potentially implicated in diabetes	78
CAS Inflammation Screen	Panel of protein targets potentially implicated in inflammation	33
CAS Liver Cancer Screen	Panel of protein targets potentially implicated in liver cancer and neoplasms	73
CAS Lung Cancer Screen	Panel of protein targets potentially implicated in lung cancer and neoplasms	66
CAS Parkinson's Disease Screen	Panel of protein targets potentially implicated in Parkinson's disease	46
CAS Prostate Cancer Screen	Panel of protein targets potentially implicated in prostate cancer and neoplasms	70

点击查看详细靶点名称列表

自訂靶點面板

Panels

+ Create New Panel

Search panels by name...

Panel Name	Description
CAS Full Pharmacology (Default)	The full panel of protein target models available through CAS to highlight potentia
Predictive_Panel_Test	

可自定义面板名称

Target	Organism	Uniprot ID
☒ Prostaglandin F2-alpha receptor	Felis catus	6L9K4
☒ Gag-pol polyprotein	Escherichia coli	A0A0A0F4Y0
☒ Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform	Rattus norvegicus	A0A0G2K344
☒ Dihydroorotate dehydrogenase (quinone)	Plasmodium falciparum Dd2	A0A0L7M6L3
☒ Bifunctional dihydrofolate reductase-thymidylate synthase	Cryptosporidium hominis	A0A054TER9
☐ Dihydrofolate reductase	Cryptosporidium hominis	A0A0W0CF37
☐ Beta-glucuronidase	Escherichia coli BL21(DE3)	A0A140N7P8
☐ Methionine aminopeptidase	Escherichia coli	A0A193LMB0
☐ Glucose-dependent insulinotropic receptor	Mesocricetus auratus	A0A1U8CD06
☐ Matrix protein 2	Influenza A virus (A/WSN/1933(H1N1))	A0A1Y0AWW2
☐ Delta-type opioid receptor	Cavia porcellus	A0A286XTF2
☐ Emopamil-binding protein-like	Cavia porcellus	A0A286XUI8
☐ 5-hydroxytryptamine receptor 2C	Cavia porcellus	A0A286XYP5
☐ 5-hydroxytryptamine receptor 1A	Cavia porcellus	A0A286Y074
☐ Thromboxane A2 receptor	Cavia porcellus	A0A286Y4I8

Create Predictive Analytics Panel

Panel Name

Predictive_Panel_Test

Description (Optional)

Cancel

Create Panel

Selected Targets

Prostaglandin F2-alpha receptor (Felis catus)

Gag-pol polyprotein (Escherichia coli)

Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform (Rattus norvegicus)

Dihydroorotate dehydrogenase (quinone) (Plasmodium falciparum Dd2)

Bifunctional dihydrofolate reductase-thymidylate synthase (Cryptosporidium hominis)

Cancel

Save

生物活性預測結果

[Return to Predictive Analytics](#)

Set Summary

Rerun Set Analysis
Complete - Last ran 09 Sep 2025

CAS Breast Cancer Screen

Total ligands: 25

Neighbors: 78

Metabolites: 53

Targets: 21

Filters

▼ pActivity

▼ Confidence Score

^ Target

☐ Androgen receptor

☐ Progesterone receptor

☐ Tumor necrosis factor

☐ Glycogen synthase kinase-3 beta

☐ Estrogen receptor

[View All](#)

Apply

▼ Organism

[Return to top](#)

Predictive_Set_09_09_2025_1002

Ligands Pharmacology

Show Structures ☒

可直接連結到相關靶點詳情頁

Ligand	Target	Organism	Known pAct	Predicted pAct	Confidence	Method
 664363-58-0	Androgen receptor	Homo sapiens	8.39	8.26	0.8	●●●●●
 664363-91-1	Progesterone receptor	Homo sapiens	-	8.02	0.39	●●●●●
 664363-58-0	Progesterone receptor	Homo sapiens	-	8.01	0.39	●●●●●
 664363-52-4	Progesterone receptor	Homo sapiens	-	7.95	0.39	●●●●●
 664363-02-4	Androgen receptor	Homo sapiens	-	7.94	0.79	●●●●●
 664363-02-4	Progesterone receptor	Homo sapiens	-	7.94	0.39	●●●●●

從左到右涉及的方法有：
MLM, SAS, SEA, SIM, XPI

使用左側篩選功能可對這些數據進行細化分析：

- 活性值(pActivity)
- 置信度分數(Confidence score)
- 靶點(Target)
- 有機體(Organism)

助於探索特定範圍內的pAct, 或有較高可信度的預測資訊。

總結

配體檢索 (Ligands)

- 基於藥物名稱、CAS RN、結構、靶點、疾病等，獲取配體資訊
- 通過類藥性、ADME、實驗參數、藥物功能、生物有機體等，進行篩選聚焦
- 查看並評估化合物的藥理學、ADME、毒理學和代謝物資訊及可視化展示

骨架檢索(Scaffolds)

- 基於結構或靶點等資訊，高效獲取相關骨架結構，並通過官能團R-Groups，進行構效關係分析
- 通過 SAR 以及MMPA，優化目標骨架或先導化合物的效力和選擇性
- 探索針對關注靶點同一結構骨架中活性最高的配體，獲取相似骨架結構與拓展

蛋白/通路/靶點與疾病檢索(Proteins & Diseases)

- 高效獲取蛋白質通路/靶點與疾病生物標誌物資訊，探索藥物作用機制和適應症

預測配體 與不同靶點的生物活性(Predictive Analytics)

- 預測特定結構針對特定靶點的藥理活性資訊
- 關注配體的脫靶效應與藥物再利用

相關資源Learn More

<https://biofinder.cas.org/>

獲取CAS BioFinder登錄訪問網址

<https://www.cas.org/solutions/biofinder-discovery-platform>

瞭解更多關於CAS BioFinder的資訊

<https://cas-biofinder.zendesk.com/hc/en-us>

獲取CAS BioFinder使用說明資訊

獲取更多技術支援與說明，請聯繫：tliu2@acs-i.org

Between problems
and progress **are**
connections that
matter

謝謝！

tliu2@acs-i.org

美国化学文摘社(CAS)

+886-975-765-358

